



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C07C 217/74, A61K 31/135, C07C 255/54, C07D 295/08, C07C 233/43, C07D 211/26, C07C 311/21, 211/60, 323/19, C07D 333/16, 213/30, 211/44, C07C 217/76, 233/25, C07D 317/54	A1	(11) International Publication Number: WO 98/38156
		(43) International Publication Date: 3 September 1998 (03.09.98)
(21) International Application Number: PCT/JP98/00780 (22) International Filing Date: 26 February 1998 (26.02.98) (30) Priority Data: 9/43940 27 February 1997 (27.02.97) JP 9/193497 18 July 1997 (18.07.97) JP (71) Applicant (for all designated States except US): TAKEDA CHEMICAL INDUSTRIES, LTD. [JP/JP]; 1-1, Doshomachi 4-chome, Chuo-ku, Osaka-shi, Osaka 541-0045 (JP). (72) Inventors; and (75) Inventors/Applicants (for US only): KATO, Kaneyoshi [JP/JP]; 2-40, Maruyamadai 2-chome, Kawanishi-shi, Hyogo 666-0152 (JP). TERAUCHI, Jun [JP/JP]; 3-5-204, Hachizuka 3-chome, Ikeda-shi, Osaka 563-0024 (JP). FUKUMOTO, Hiroaki [JP/JP]; 18-D74-107, Tsukumodai 5-chome, Suita-shi, Osaka 565-0862 (JP). KAKIHANA, Mitsuru [JP/JP]; 4-2, Tsukushigaoka 9-chome, Kita-ku, Kobe-shi, Hyogo 651-1212 (JP).		(74) Agents: ASAHINA, Tadao et al.; Osaka Plant of Takeda Chemical Industries, Ltd., 17-85, Jusohonmachi 2-chome, Yodogawa-ku, Osaka-shi, Osaka 532-0024 (JP). (81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, GW, HU, ID, IL, IS, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: AMINE COMPOUNDS, THEIR PRODUCTION AND USE AS AMYLOID-BETA PRODUCTION INHIBITORS		
<div style="text-align: center;"> </div> <div style="text-align: right;"> (I) </div>		
(57) Abstract <p>A compound of formula (I) wherein Ar is an aromatic ring assembly group which may be substituted or a fused aromatic group which may be substituted; X is (i) a bond, (ii) -S-, -SO- or -SO₂-, (iii) C₁₋₆ alkylene, C₂₋₆ alkenylene or C₂₋₆ alkynylene, etc., (iv) -CO-O- or (v) -(CH₂)_p-X¹-(CH₂)_q-, -(CH₂)_r-CO-X¹-, -SO₂-NR⁸- or -(CH₂)_r-SO₂-NR⁸- wherein X¹ is O or NR⁸, R⁸ is H, a hydrocarbon group which may be substituted or an acyl, p is 0 to 5, q is 1 to 5, p+q is 1 to 5, and r is 1 to 4; Y is a divalent C₁₋₆ aliphatic hydrocarbon group optionally containing O or S, which may be substituted; R¹ and R² each is H or a lower alkyl which may be substituted, or R¹ and R² form an N-containing heterocyclic ring which may be substituted; Ring A is a benzene ring which may be further substituted; and Ring B is a 4- to 8-membered ring which may be further substituted, or a salt thereof has the effect of inhibiting amyloid-β protein production and/or secretion and is useful as a pharmaceutical composition for preventing and/or treating the neurodegenerative disease, etc.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

DESCRIPTION

AMINE COMPOUNDS, THEIR PRODUCTION AND USE AS AMYLOID-BETA PRODUCTION INHIBITORS

5 TECHNICAL FIELD

The present invention relates to an amine compound having an excellent effect of inhibiting production and/or secretion of amyloid- β protein, a production and use thereof. Especially, it is effective for
10 preventing and/or treating, for example, neurodegenerative diseases, amyloid angiopathy, neurological disorders caused by cerebrovascular disorders, and so forth.

15 BACKGROUND ART

Alzheimer's disease is a neurodegenerative disease, which is characterized by the degeneration and loss of neuronal cells accompanied by the formation of senile plaques and neurofibrillary tangles. Senile plaque
20 that is the most characteristic in Alzheimer's disease consist of essentially amyloid- β protein (hereinafter referred to as A β) [see Biochem. Biophys. Res. Commun., 122, 1311 (1984)] and other intracerebral components. It is known that A β comprised of 40 or 42 amino acids
25 (hereinafter referred to as A β ₁₋₄₀ and A β ₁₋₄₂, respectively) is toxic to neurons and induces neurofibrillary changes.

Some patients with familial Alzheimer's disease are known to have APP (amyloid precursor protein) gene mutation, and it is well known that the cells
30 transfected with such mutated gene produce and secrete an increased amount of A β [for example, see Nature, 360, 672 (1992); Science, 259, 514 (1993); Science, 264, 1336 (1994), etc.].


Based on the information, medicines which inhibit production and/or secretion are useful for preventing and/or treating diseases caused by A β (e.g., Alzheimer's disease, Down's syndrome, etc).

Alternatively, secreted form of amyloid precursor protein (sAPP) is reported to have neurotrophic factor like property (Neuron, 10, 243-254, 1993). As neurotrophic factor like property, 1) survival and preserving effect to the neuronal cell; 2) stimulating the synapse formation; 3) protection of neuronal cell death; and 4) long term potentiation in hippocampus are given as examples. By the above-mentioned property, drugs which stimulate the sAPP secretion are also useful in preventing and treating 1) neurodegenerative diseases such as dementia (e.g., senile dementia, amnesia, etc.), Alzheimer's disease, Down's syndrome, Parkinson's disease, Creutzfeldt-Jacob disease, amyotrophic sclerosis on lateral fasciculus, Huntington's disease, multiple sclerosis, etc., 2) neurological disorders involved in cerebrovascular disorders (e.g., cerebral infarction, encephalorrhagia, etc.), a head injury or an injury of spinal cord, and so forth.

EP-A-652009 discloses peptide derivatives which is a protease inhibitor exhibiting an A β production inhibiting effect in *in vitro* experiments using cell lines.

On the other hand, the following bicyclic amine compounds are known.


1) JP-A-2-96552 (USP 5,137,901) discloses a compound of the formula:



wherein Y represents a straight-chain or branched,


substituted or unsubstituted alkylene chain having up to 6 carbon atoms; Z represents a group of the formula: -NR²R³, -OR⁴, or the like; R² and R³ are identical or different and represent hydrogen, alkyl, alkenyl or cycloalkyl, or represent aryl which may be substituted by halogen, etc.; R⁴ represents hydrogen, alkyl, alkenyl, or the like; R¹ represents hydrogen, alkyl, aralkyl, heteroarylalkyl or a group of the formula: -(Y¹-Z¹) in which Y¹ and Z¹ are identical or different and have the same meanings as Y and Z; A and D each represents a group of the formula: -CH₂, O, S or NR¹³, or the moiety of -CH or N of a double bond C=C or C=NH, with the proviso that either only A or only D represents oxygen, sulfur or N-R¹³; R¹³ represents hydrogen, alkyl, alkoxy, acyl, alkoxycarbonyl or alkylsulfonyl; B represents a group of the formula: -CH₂ or $\equiv\text{CH}$ or the moiety of -CH or N of a double bond C=C or C=N; C represents a group of the formula: $\equiv\text{CH}$, or the moiety of C of a double bond C=C or C=N; E and F are identical or different and each represents hydrogen, alkyl, alkoxy, halogen, nitro, cyano, trifluoromethyl, trifluoromethoxy or a group of the formula: -CONR²R³ in which R² and R³ have the same meanings as above, or E and F together form a substituted or unsubstituted carbocycle having 6 carbon atoms, which is agonist, partial agonist and antagonist on the serotonin receptors and is suitable for the treatment of central nervous system disorders, etc.

2) JP-A-63-77842 discloses a compound of the formula:




5

25




30

4) WO 95/32967 discloses amide derivatives of the formula:



wherein A is CONR where R is hydrogen or C₁₋₆ alkyl; Q is an optionally substituted 5 to 7-membered heterocyclic ring containing 1 to 3 heteroatoms selected from oxygen, nitrogen or sulphur; R₁ is hydrogen, halogen, etc.; R₂ and R₃ are independently hydrogen, halogen, etc.; R₄ and R₅ are independently hydrogen or C₁₋₆ alkyl; R₆ is halogen, hydroxy, etc.; R₇ and R₈ are independently hydrogen, C₁₋₆ alkyl, aralkyl, or together with the nitrogen atom to which they are attached from an optionally substituted 5 to 7-membered heterocyclic ring containing one or two heteroatoms selected from oxygen, nitrogen or sulphur; m is 0 to 4; and n is 0, 1 or 2, which has 5HT_{1D} receptor antagonist activity and is useful for the treatment of various CNS disorders.

5) EP-A-754455 discloses a pharmaceutical compositions for the therapeutic application as neuroprotectors in Parkinson's and Alzheimer's diseases containing a compound of the formula:




wherein X is H, halogen, alkoxy, alkyl, alkylthio, aryl, aryloxy; R is H, CH₃, or other aliphatic, alicyclic or aryl radicals; R' is H, CH₃, or other aliphatic or alicyclic C₁-C₃ radicals, or an aryl or arylalkyl, or a radical the same as those indicated for R''; and R'' is H, CH₃, or other aliphatic or alicyclic C₁-C₃ radicals, or an aryl or arylalkyl, or an acetylene or allene

group, being potent selective monoamine oxydase B inhibitors.

The conventional A β production inhibitors for the treatment of Alzheimer's disease are problematic in their oral absorbability, stability, etc. and are therefore unsatisfactory as medicines. It is desired to develop a compound which is different from the known compounds mentioned above in its chemical structure and which have an excellent inhibitory effect on A β production and/or secretion and is therefore satisfactorily used in medicines.

DISCLOSURE OF INVENTION

We, the present inventors have studied various compounds having an inhibitory effect on A β production and/or secretion and, as a result, have succeeded in, for the first time, the production of novel a compound of the formula:



wherein Ar represents an aromatic ring assembly group which may be substituted or a fused aromatic group which may be substituted;
X represents (i) a bond, (ii) -S-, -SO- or -SO₂-, (iii) a C₁₋₆ alkylene, C₂₋₆ alkenylene or C₂₋₆ alkynylene group, each of which may be substituted by 1 to 3 substituents selected from the group consisting of oxo and C₁₋₆ alkyl, (iv) -CO-O- or (v) a group of the formula: -(CH₂)_p-X¹-, -(CH₂)_p-X¹-(CH₂)_q-, -(CH₂)_r-CO-X¹-, -SO₂-NR⁸- or -(CH₂)_r-SO₂-NR⁸- wherein X¹ represents O (oxygen atom) or NR⁸, R⁸ represents a hydrogen atom, a hydrocarbon group which may be substituted or an acyl, p represents an integer of 0 to 5, q represents an integer of 1 to 5,

p+q is an integer of 1 to 5, and r represents an integer of 1 to 4;

Y represents a divalent C₁₋₆ aliphatic hydrocarbon group which may contain an oxygen atom or a sulfur atom and may be substituted;


R¹ and R² each represents a hydrogen atom or a lower alkyl which may be substituted, or

R¹ and R² form, taken together with the adjacent nitrogen atom, a nitrogen-containing heterocyclic ring which may be substituted;

Ring A represents a benzene ring which may be further substituted apart from the group of the formula: -X-Ar wherein each symbol is as defined above; and

Ring B represents a 4- to 8-membered ring which may be further substituted apart from the group of the formula: -Y-NR¹R² wherein each symbol is as defined above;

provided that, when the fused ring to be formed by Ring A and Ring B is an indole ring, the group of the formula: -X-Ar wherein each symbol is as defined above is substituted on 4-, 6- or 7-position of the indole ring, or a salt thereof [hereinafter sometimes referred to as compound (I)], which is characterized by the chemical structure in that the benzene ring A of the skeleton of the formula:



wherein the symbols have the same meanings as above, is substituted by the group of the formula: -X-Ar wherein the symbols have the same meanings as above. We have found for the first time that compound (I), being based on its specific chemical structure, has an unexpected, excellent inhibitory effect on A β production and/or secretion, that a compound of the formula: